



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Cerebral organoids as tools to identify the developmental roots of autism

Citation for published version:

Chan, WK, Griffiths, R, Price, DJ & Mason, JO 2020, 'Cerebral organoids as tools to identify the developmental roots of autism', *Molecular Autism*, vol. 11, no. 1, pp. 58. <https://doi.org/10.1186/s13229-020-00360-3>

Digital Object Identifier (DOI):

[10.1186/s13229-020-00360-3](https://doi.org/10.1186/s13229-020-00360-3)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Molecular Autism

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Cerebral organoids as tools to identify the developmental roots of autism

Wai Kit Chan[¶], Rosie Griffiths[¶], David J. Price and John O. Mason*

Centre for Discovery Brain Sciences and Simons Initiative for the Developing Brain, University of Edinburgh, George Square, Edinburgh EH8 9XD

[¶]These authors contributed equally

*Corresponding author: John Mason John.Mason@ed.ac.uk

Keywords: Autism Spectrum Disorder; Cerebral organoids; Embryonic brain development.

List of abbreviations: AD: Alzheimer's Disease, AP: action potential; AS: Angelman Syndrome, ASD: Autism Spectrum Disorder; CP: cortical plate; FXS: Fragile X Syndrome; GW: gestational week; GWAS: genome wide association study; IPC: intermediate progenitor cell; iPSC: induced pluripotent stem cell; ISVZ: inner subventricular zone; IZ: intermediate zone; LTCC: L-type calcium channel; MZ: marginal zone; OPC: oligodendrocyte progenitor cell; oRG: outer radial glia; osmFISH: ouroboros single-molecule fluorescence in situ hybridization; oSVZ: outer subventricular zone; p.c.w.: post-coital week; PMS: Phelan-McDermid Syndrome; PSC: pluripotent stem cell; RGC: radial glial cell; scRNA seq: single cell RNA sequencing; SFARI: Simons Foundation for Autism Research Initiative; SP: subplate; SVZ: subventricular zone; TS: Timothy Syndrome; VZ: Ventricular zone;

Abstract

Some Autism Spectrum Disorders (ASD) likely arise as a result of abnormalities during early embryonic development of the brain. Studying human embryonic brain development directly is challenging, mainly due to ethical and practical constraints. However, the recent development of cerebral organoids provides a powerful tool for studying both normal human embryonic brain development and, potentially, the origins of neurodevelopmental disorders including ASD.

Substantial evidence now indicates that cerebral organoids can mimic normal embryonic brain development and neural cells found in organoids closely resemble their in vivo counterparts. However, with prolonged culture, significant differences begin to arise. We suggest that cerebral organoids, in their current form, are most suitable to model earlier neurodevelopmental events and processes such as neurogenesis and cortical lamination. Processes implicated in ASDs which occur at later stages of development, such as synaptogenesis and neural circuit formation, may also be modelled using organoids. The accuracy of such models will benefit from continuous improvements to protocols for organoid differentiation.

1. Introduction

Autism spectrum disorders (ASDs) are a group of neurodevelopmental disorders that affect as many as 1 in 59 children (CDC, 2012). They are characterized by impairments in social interaction and communication and repetitive and restricted patterns of behaviour, interests or activities. While these symptoms can be found in any individual across the spectrum, the severity of the symptoms presented varies, ranging from very mild to very severe. Individuals with ASD may also present distinct combinations of comorbid features and diagnoses that are not part of the disorder they were diagnosed with, such as gastrointestinal symptoms, epilepsy, sleep disruptions, or motor disturbances. This clinical heterogeneity makes it difficult to find a unifying biological hypothesis to address all the features of ASD and the underlying genetic causes of ASDs are still under debate.

Studies of large family cohorts have identified at least 65 ASD risk genes with high confidence (Fischbach and Lord, 2010; Zhao et al., 2007) and hundreds more candidate genes. However, this only accounts for about 30% of ASD cases, the remainder having nonsyndromic idiopathic ASD in which the cause is unknown (Fernandez and Scherer, 2017). It is estimated that as many as 300 – 1000 genes could be targets for rare mutations which greatly increase the risk of ASD, potentially explaining some idiopathic cases (He et al., 2013; Ronemus et al., 2014). This extreme genetic heterogeneity makes it very difficult to map the relationship between genotype and phenotype in ASD (De Rubeis and Buxbaum, 2015). However recent work using network approaches suggests that autism risk genes converge on a small number of biological pathways and processes (de la Torre-Ubieta et al., 2016; Gilman et al., 2011; Mullins et al., 2016; Wen et al., 2016). Gene set enrichment analyses have shown that genes associated with ASD converge

on pathways and processes that contribute to embryonic brain development, including chromatin remodelling, neurogenesis and cortical lamination; neuronal physiological maintenance; and synaptic processes. The convergence of many ASD-risk genes on common molecular pathways may help explain how a genetically heterogeneous population of individuals exhibit similar symptoms.

One such point of convergence is synaptogenesis and synapse physiology (Bhandari et al., 2020; de la Torre-Ubieta et al., 2016; Guang et al., 2018). One of the earliest ASD-risk genes identified, *SHANK3*, and many other ASD-risk genes identified subsequently are directly involved in synapse physiology, highlighting the clear role that dysregulation of synaptogenesis and synaptic transmission play in ASD pathophysiology. So, ASD has often been viewed as a disorder of synaptic dysfunction (Zoghbi and Bear, 2012). However, many neurodevelopmental processes which occur during embryonic and fetal stages, such as neurogenesis and cortical lamination, are also a point of convergence for ASD risk genes, indicating that ASD can arise from an earlier point in development (Casanova, 2014; Packer, 2016; Vaccarino et al., 2009). This view is supported by a recent genome wide association study (GWAS) which showed that a large proportion of ASD risk genes analysed were expressed most highly during fetal corticogenesis (Grove et al., 2019). These early stages of brain development are highly dynamic. One could hypothesize that small changes during these processes could lead to larger effects later. Key stages of brain development at which ASD-related genes may act are described in Box1.

Studying human prenatal brain development directly remains a major challenge, due to scarcity of material and ethical constraints on research using human embryos. Much of our present understanding of brain development is therefore based on studies using model organisms, primarily the mouse. However, there are important differences between mouse and human brain development (Florio and Huttner, 2014). The recent advent of cerebral organoids offers the potential to study human brain development directly. Cerebral organoids are 3D cultured cell aggregates derived from pluripotent stem cells (PSCs) which closely resemble embryonic brain tissue. They contain many of the cell types found in embryonic brains, locally organized in a similar way to that found *in vivo*, and exhibit similar behaviours (Lancaster et al., 2013), but the spatial organisation along major axes (anteroposterior, dorsoventral and mediolateral) found in embryos is absent in organoids. Organoids have the potential to be invaluable tools for studying

both normal development and the developmental origins of neurodevelopmental disorders including ASDs. A number of studies have already used organoids to model neurodevelopmental disorders as summarised in Table 1.

Protocols for generating human cerebral organoids fall into two main categories. In the first, PSCs are aggregated and allowed to differentiate in the absence of any specific added differentiation cues. Such protocols, exemplified by Lancaster et al., 2013, produce heterogeneous organoids containing areas corresponding to various regions of the brain, such as dorsal and ventral forebrain, hindbrain, hippocampus or choroid plexus. Alternatively, many protocols include the addition of specific cues that promote formation of a specific region of the brain, such as dorsal forebrain, ventral forebrain, midbrain, hypothalamus or thalamus (Birey et al., 2017; Qian et al., 2016; Xiang et al., 2019). Such regionalized organoids can be used to investigate developmental patterning and the effects of mutations on individual brain regions. A recent study summarized all widely used organoid protocols and compared the transcriptomic profiles of the organoids grown using a range of published protocols. They found that while each protocol produced organoids with similar cellular composition, the differentiation trajectories differed between protocols (Tanaka et al., 2020). Before using cerebral organoids to investigate normal development or disease states, it is important to understand how accurately they can recapitulate the *in vivo* system and what their limitations are. Comparing the transcriptomes of cortical organoids with those of human fetal cortex shows encouraging similarities. At a global level, the transcriptome of dorsal forebrain organoids grown for 40-100 days correlated best with fetal cortex tissue at ages 8-16 p.c.w, indicating that organoids develop at a similar rate to the fetus (Amiri et al., 2018; Luo et al., 2016; Mariani et al., 2015). The epigenome of organoids is also similar to that of fetal tissue - analysis of histone modifications showed cortical organoids were more like fetal brain tissue than adult brain tissue or pluripotent stem cells (Amiri et al., 2018).

Single cell analysis of telencephalic organoids and fetal human cortex showed that they contain very similar cell types. While excitatory neurons are the most numerous cell type within the cortex, several other vital cell types are also present. Fetal cortex at ages 15-23 p.c.w contains, in decreasing order of proportion: excitatory neurons, inhibitory neurons that migrated from the ventral telencephalon, radial glial, astrocytes, microglial, with a small subset of glia, endothelial cells, oligodendrocyte progenitor cells and Cajal-Retzius cells (Amiri et al., 2018; Fan et al.,

2018; Zhong et al., 2018). At early stages (1-3 months in culture), human dorsal forebrain organoids contain mostly excitatory neurons, radial glial and intermediate progenitor cells (Amiri et al., 2018; Birey et al., 2017; Velasco et al., 2019; Yoon et al., 2019), while at six months there is an increase in astroglia and inhibitory interneurons (possibly of the olfactory bulbs) begin to appear (Velasco et al., 2019). Ventral patterned cortical organoids on the other hand, contain GABAergic neurons, ventral progenitors and astroglial cells, the key cell types found in the cognate brain structures *in vivo*. Although organoids show similar cell type composition to that found *in vivo*, a recent large scRNA-seq analysis of organoids and primary tissues, showed organoids lacked cell type and sub-type fidelity indicated by co-expression of different cell-type markers in organoids when compared to fetal brain (Bhaduri et al., 2020). Organoid transcriptomes change during differentiation in accordance with the development of the fetal cortex. Modules of genes co-expressed during cortex development *in vivo* are conserved in organoids. These include upregulated genes associated with synaptic transmission, cell adhesion and neuron differentiation and downregulated cell cycle genes (Amiri et al., 2018; Luo et al., 2016). Notably, several gene modules co-expressed during organoid development are enriched with SFARI genes, a curated list of genes associated with ASD, and a quarter of SFARI genes are differentially expressed during organoid differentiation (Amiri et al., 2018). *In vivo*, genes linked to ASD and intellectual disability, including genes associated with chromatin remodeling, Wnt and Notch signalling, were most highly expressed at 8-16 p.c.w. (Hormozdiari et al., 2015). Another single-cell transcriptome analysis found around 84% of genes mutated in disorders affecting neurogenesis showed the same developmental expression trajectory in organoid and fetal cells (Camp et al., 2015). These studies suggest that organoids can recapitulate the timeline of events during normal human development during which ASD causative effects could be taking place.

2. Possible developmental origins of ASD

2.1 Abnormal neurogenesis and growth of the cerebral cortex

One obvious difference between mouse and human brains is the disproportionately increased size of the human cerebral cortex. The mechanisms that led to the dramatic expansion of the human cortex are outlined in Box 2. Enlarged head size is a common feature of ASDs. Some 14%–34% of autistic patients show macrocephaly (Sacco et al., 2015), due to increased surface area rather

than increased cortical thickness (Ohta et al., 2016). Increased brain volume has been linked to the emergence and severity of autistic social deficits (Hazlett et al., 2017). Analysis of head circumference of children with ASD over the first year of life showed an accelerated increase in head size (Courchesne et al., 2003). Aberrant brain growth could be due to changes to the balance between proliferation and differentiation of neural progenitor cells in the embryo (Courchesne et al., 2011; Florio and Huttner, 2014). Cell-cycle genes have been implicated in ASD based on differential expression in postmortem ASD brains (Chow et al., 2012). A systems biology approach analyzing total brain volume and gene expression levels (in blood, given the unavailability of brain tissue) in ASD toddlers also implicated cell-cycle genes in regulation of brain size (Pramparo et al., 2015). Mutations in cell cycle control genes have been found in ASD patients. For example, mutations in the transcriptional regulator *ANKRD11* (Ankyrin repeat domain 11) contribute to ASD (Iossifov et al., 2014; Marshall et al., 2008). *ANKRD11* regulates neural progenitor proliferation through interaction with histone deacetylases (Zhang et al., 2004). Mutations in *PTEN* (phosphatase and tensin homologue on chromosome ten), a phosphatase that acts to inhibit the AKT/mTOR pathway, are associated with macrocephaly (Butler et al., 2005) and ASD (Conti et al., 2012; De Rubeis et al., 2014). *Pten* heterozygous mutant mice have an increased number of neural progenitors and a decreased total cell number in the mature cerebral cortex, due to fewer neural progenitors exiting the cell cycle (Chen et al., 2015). In contrast, *PTEN* homozygous mutants in human cerebral organoids exhibited an expanded VZ and oSVZ, delayed neuronal differentiation and surface expansion and folding (Li et al., 2017). Numerous mouse studies demonstrate that mutations in cell cycle genes can cause overgrowth of cerebral cortex. For example, knockdown of *Ankrd11* by *in utero* electroporation at E13/14 revealed decreased neural progenitors and underproduction of neurons, an effect that was rescued by administering a histone acetyltransferase inhibitor (Gallagher et al., 2015). Overexpression of Cdk4/Cyclin D1 via *in utero* electroporation of mouse brains inhibited the switch from proliferation to differentiation, resulting in a cortex with a larger surface area, reminiscent of the increased surface area found in human ASD patients (Lange et al., 2009; Pilaz et al., 2016).

2.1.1 Progenitor proliferation and neurogenesis in human cerebral organoids

Human cerebral organoids faithfully recapitulate embryonic cortical structures – they generate neuroepithelium and PAX6⁺ SOX2⁺ progenitors are found at its apical surface, closely

resembling the embryonic ventricular zone. An adjacent layer of TBR2⁺ intermediate progenitor cells indicates the presence of an SVZ, and expression of neuronal markers TBR1 and MAP2 at the basal surface indicate cortical plate (Lancaster et al., 2013). The presence of an outer SVZ (oSVZ) as indicated by expression of outer RG markers HOPX and PTPRZ1 has also been reported in some cerebral organoids (Qian et al., 2016; Qian et al, 2020). This is important, as oSVZ progenitors are thought to drive cortical expansion in human and are absent in mice. Thus, human organoids are likely to be an invaluable tool to study this population of cells.

Human cortical organoids have been used to study disorders with a brain size abnormality. Organoids grown from iPSCs derived from a patient with severe microcephaly had reduced neural tissue, resembling the patient phenotype. Analysis of the early stages of organoid differentiation showed they had smaller neuroepithelia, fewer radial glial cells and more neurons indicating an imbalance of the symmetric and asymmetric divisions of neural progenitor cells (Lancaster et al., 2013).

In a study of idiopathic ASD, iPSCs were derived from family members with an ASD and unaffected close relatives, then cerebral organoids were grown from each. The transcription factor *FOXP1* was found to be significantly overexpressed in the ASD patient-derived organoids, driving an accelerated cell cycle (Mariani et al., 2015). Transcriptome analysis showed increased expression of genes associated with neural differentiation and synaptic transmission in ASD-organoids (Mariani et al., 2015), both of which have been linked to ASD (De Rubeis et al., 2014; Gilman et al., 2011; Pinto et al., 2014). ASD-organoids showed increased neural maturation and surplus GABAergic neurons but no effect on excitatory neuron number, indicating an imbalance of excitatory/inhibitory neurons, a phenotype believed to underlie some cases of autism. Inhibition of *FOXP1* expression restored GABAergic neuronal numbers to normal (Mariani et al., 2015).

CHD8, one of the most commonly mutated genes in ASD, can negatively regulate WNT signalling, an essential signalling pathway in brain development. Transcriptome analysis of forebrain organoids derived from *CHD8*^{+/-} iPSC lines and their isogenic controls showed dysregulation of genes associated with neurogenesis, WNT signalling and ECM components (Wang et al., 2017). Notably, there was significant overlap of these differentially expressed genes when compared to those found in NPCs and neurons derived from *CHD8*^{+/-} iPSCs compared to controls in 2D culture experiments (Wang et al., 2015). Similarly, 23% of

differentially expressed genes found in *CHD8*^{+/-} organoids were also found in idiopathic ASD organoids, with these overlapping genes being enriched with neurogenesis associated genes (Mariani et al., 2015; Wang et al., 2017).

2.2 Defective neuronal migration / cortical lamination

Disruption to cortical lamination may be a common feature of brain development in ASD (Casanova, 2014; Stoner et al., 2014). Cortical layers form progressively during embryonic development, with deep layer neurons being born first and later-born neurons migrating past them to form the characteristic six-layered laminar architecture of the cortex. Defects in migration could be indirect effects of altered cell cycle dynamics or proliferation as migration defects are also observed in the mouse models of the genes discussed above. For example, in *Ankrd11* mutant mice more cells are retained in the VZ and SVZ, resulting in fewer cells in the cortical plate. Furthermore, there were fewer Satb2-expressing superficial layer neurons and Tbr1⁺ deep-layer neurons were positioned inappropriately (Sirmaci et al., 2011). Pten heterozygous mutant mice showed an increase in superficial layer Cux1-expressing neurons (Chen et al., 2015).

The transcription factor *TBR1* is required for normal cortical lamination and has been implicated in ASD (Bedogni et al., 2010; De Rubeis et al., 2014; Hevner et al., 2001). Tbr1 is expressed in deep layer neurons (layer 5/6) and in *Tbr1*^{-/-} mutant mice, neurons in layer 5 and layer 6 of the cortical plate are mixed and there is no clear distinction between them (Bedogni et al., 2010; Hevner et al., 2001). Tbr1 is required to maintain layer 6 identity in the postnatal cortex - specific deletion of *Tbr1* in layer 6 led to increased expression of regulators of layer 5 identity such as *Fezf2* and *Bcl11b* and a decrease in layer 6 markers/regulators *Foxp2* and *Tle4* (Fazel Darbandi et al., 2018).

Before initiating radial migration, newborn cortical neurons undergo a multipolar to bipolar morphology change. This transition is disrupted in mice deficient for a number of genes implicated in ASD including *Foxg1* (Miyoshi and Fishell, 2012) and *Fmr1* (La Fata et al., 2014). *Foxg1*, a transcription factor, is expressed in neural progenitors and has multiple roles in forebrain development in mice (Manuel et al., 2011; Shen et al., 2019). Downregulation of Foxg1 expression is required to allow cells to progress from multipolar to bipolar morphology before migrating into the cortical plate (Miyoshi and Fishell, 2012). Delay in multipolar to bipolar transition impairs the coordinated integration of excitatory neurons with inhibitory

interneurons, ultimately affecting the ratio of excitatory to inhibitory neurons in the developing cortex (Mariani et al., 2015; Miyoshi and Fishell, 2012).

2.2.1 Neuronal migration and cortical lamination in cerebral organoids

Forebrain organoids contain radially aligned RGC processes as seen *in vivo* (Qian et al., 2018). Live imaging of neurons migrating from organoids onto a Matrigel surface showed similar migration rates to ferret cortical explants (Bershteyn et al., 2017). Using this model, migration defects were observed in organoids derived from iPSCs from Miller-Dieker Syndrome (MDS) patients; a severe cortical malformation disorder caused by defective cortical neuronal migration. Imaging analysis showed fewer neurons migrating out of the organoid, reduced migration speed and track straightness (Bershteyn et al., 2017).

Migratory defects can also be examined by using an ‘assembloid’ approach. By co-culturing ventral telencephalic organoids with dorsal cortical organoids, interneurons from the ventral organoids were shown to migrate towards the dorsal forebrain as observed *in vivo* (Birey et al., 2017). Using this model, assembloids were generated using hiPSCs derived from patients with Timothy Syndrome (TS), a neurodevelopmental disorder characterized by ASD and epilepsy (Birey et al., 2017). These assembloids exhibited migration defects in interneurons and increased residual calcium following depolarization in TS neurons. Timothy syndrome is caused by a gain-of-function mutation in an L-type calcium channel (LTCC) subunit and incubating organoids with an LTCC blocker successfully reversed the migratory defects (Birey et al., 2017).

Human cortical organoids show some lamination (Figure 1), with progenitor cells located towards the central lumen and differentiated neurons located towards the outside. Expression of TBR1, CTIP2 and SATB2 showing similar lamination in day 84 organoids as found in neocortex at 23 GWs (Qian et al., 2016; Saito et al., 2011). A recent study showed that cutting organoids into thick organotypic slices, thereby improving nutrient access from the culture medium, greatly enhanced lamination as indicated by larger and more distinct oSVZ and CP layers. Using this method, they identified lamination defects in organoids with a DISC1 mutation, which has been associated with schizophrenia and autism (Kamiya et al., 2005; Qian et al., 2020).

2.3 Abnormal synaptogenesis in ASD

Following neurogenesis, neuronal migration and cortical lamination, neurons next make connections with their appropriate synaptic partners, thus beginning the formation of neural circuits. This is known to be important in the pathophysiology of ASD as synaptogenesis is another point of convergence for ASD-risk genes (Courchesne et al., 2020; de la Torre-Ubieta et al., 2016). Here, we discuss some of the well-known ASD-risk genes involved in synaptogenesis, but this is by no means an exhaustive list. A more comprehensive list is provided in a recent review by Guang and colleagues (Guang et al., 2018).

Some of the first genes implicated in ASD, such as the multiple Ankyrin repeat domain 3 gene SHANK3, are directly involved in synapse formation and function. SHANK3 was found to be disrupted in a child affected with Phelan-McDermid syndrome (PMS) in which patients show poor eye contact, global developmental delay, decreased socialization and stereotypic movements (Bonaglia et al., 2001). Analysis of more than 60 additional patients showed that heterozygous loss of SHANK3 is responsible for the neurological phenotypes (Anderlid et al., 2002; Dhar et al., 2010). Subsequently, SHANK3 mutations have been found in numerous ASD patients (Durand et al., 2007; Gauthier et al., 2010; Moessner et al., 2007). The SHANK protein family comprises of SHANK1, 2 and 3, scaffold proteins which localize to synapses and interact with components of the postsynaptic density including guanylate kinase-associated protein (PSD95), Homer, cortactin-binding protein, and the somatostatin receptor and act to stabilize the PSD-95/Shank/Homer complexes at the postsynaptic density (Lim et al., 1999).

In mice, Shank1 is required for correct synapse maturation and function. Deleting Shank1 leads to smaller dendritic spines, thin postsynaptic densities and ultimately weakened synaptic transmission. Mutant mice showed impaired contextual fear memory, poor long-term retention of a spatial task and anxiety-like behaviour (Hung et al., 2008). Shank1's role in synapse maturation and function was corroborated in *in vitro* studies where Shank1 was overexpressed in hippocampal neurons, leading to increased maturation and size of dendritic spines (Sala et al., 2001). Mice heterozygous for Shank3 showed reduced basal synaptic transmission in hippocampal CA1 neurons and decreased long-term potentiation, an important mechanism in retaining nascent synapses (Bozdagi et al., 2010). Deleting Shank3B caused reduction in postsynaptic proteins such as Homer and PSD93. The thickness and length of the postsynaptic densities were also reduced in addition to lowered spine density. This was accompanied by a reduction in excitatory synaptic transmission in the striatum. The mice displayed anxiety-like

behaviours and decreased social interactions similar to human PMS patients (Peca et al., 2011). *In vitro* studies knocking down Shank3 in hippocampal neurons led to a lower number but increased length of dendritic spines supporting evidence from the mouse model that Shank3 is crucial for dendritic spine function (Roussignol et al., 2005). Furthermore, expressing Shank3 in aspiny cerebellar granule neurons was found to be sufficient to induce functional dendritic spines (Roussignol et al., 2005). Recent studies using human iPSCs with SHANK3 knockdown also found defects in excitatory and inhibitory synaptic transmission (Huang et al., 2019).

Fragile X syndrome (FXS) is an ASD caused by insufficient expression of the *FMR1* gene. Symptoms include intellectual disability, motor abnormalities, anxiety, speech delay, gaze avoidance and stereotyped repetitive behaviours (Hagerman et al., 1984). Postmortem neuropathological studies on FXS patients revealed structural defects of dendritic spines (Comery et al., 1997; Irwin et al., 2000). *FMR1* encodes FMRP, a multi-functional mRNA binding protein involved in the transport and localization of a subset of dendritic mRNAs (Bagni and Greenough, 2005; Laggerbauer et al., 2001; Li et al., 2001). FMRP is enriched in neurons and especially at the dendrites, where it represses the translation of many mRNAs that play important roles in synapse formation and synaptic plasticity (Brown et al., 2001; Darnell et al., 2001; Feng et al., 1997; Miyashiro et al., 2003; Weiler et al., 1997). FMRP represses the translation of many mRNAs (O'Donnell and Warren, 2002) leading to an increase in rate of basal protein synthesis in the hippocampus of *Fmr1* null mutant mice (Osterweil et al., 2010; Qin et al., 2005).

Rett syndrome is an ASD in which patients appear to develop normally up to 6 – 18 months of age but then head growth decelerates, leading to microcephaly by the second or third year of life. Other symptoms include loss of language, social withdrawal, lack of eye contact, lack of response to social cues and stereotypic hand movements (Chahrour and Zoghbi, 2007). Most cases of Rett Syndrome are caused by mutations in the *MECP2* gene, which encodes Methyl-CpG-binding protein 2, a nuclear protein that binds to methylated 5-hydroxymethylcytosine or CpG sites required for chromatin organization and transcriptional regulation (Hendrich and Bird, 1998; Lewis et al., 1992; Mellen et al., 2012). *MECP2* expression is most abundant in neurons with the expression level increasing postnatally as neurons mature (Shahbazian et al., 2002). *MECP2* regulates expression of Brain-derived neurotrophic factor (BDNF) which is a critical synaptic maturation factor maturation (Chahrour et al., 2008; Jordan et al., 2007; Nuber et al.,

2005; Tudor et al., 2002). MECP2 is important for the development and the maintenance of synapses. It is essential for the transcription of biosynthetic enzymes crucial for neurotransmitters in respective neurons such as tyrosine hydroxylase in catecholamine neurons, GAD in inhibitory neurons and neuropeptides important for neuronal physiology such as corticotropin-releasing hormone, BDNF and somatostatin (Chao et al., 2010; Samaco et al., 2009). Rodent models lacking functional MeCP2 reproduce features of Rett syndrome patients (Chen et al., 2001; Patterson et al., 2016; Pelka et al., 2006; Wu et al., 2016). Mouse models also showed a decrease in synaptic density and reduced LTP and synaptic plasticity (Moretti et al., 2006). Similarly, *in vitro* data from neurons derived from human iPSC of Rett syndrome patients revealed decreased spontaneous postsynaptic currents with fewer synapses (Marchetto et al., 2010).

The finding that MECP2 overexpression or underexpression leads to Rett syndrome-like phenotypes further complicates the role of MECP2 suggesting that gene dosage is an important factor in ASD. This also suggests that ASD might reflect a failure of homeostatic regulation of synaptic function which makes sense as optimal synaptic function only occurs within a narrow dynamic range. Too much or too little of a protein might tilt the balance out of this range, resulting in ASD.

3. Organoids generated by current protocols are not mature enough for a thorough investigation of their electrical properties

Neuronal activity has been found in human organoids older than 3 months.

Immunocytochemistry shows co-localization of pre- and post-synaptic markers in cortical organoids suggesting the formation of synapses (Birey et al., 2017; Pasca et al., 2015; Velasco et al., 2019; Xiang et al., 2019). Action potentials have been recorded in 50-80% of neurons within organoids in response to depolarization (Birey et al., 2017; Pasca et al., 2015; Xiang et al., 2019) and spontaneous firing has also been observed in organoids, which is lost after incubation with neurotransmitter antagonists (Birey et al., 2017; Pasca et al., 2015; Trujillo et al., 2019; Xiang et al., 2019). Furthermore, deep layer neurons in fetal neocortex show complex dendrite morphology at GW26 (Zhong et al., 2018) and neurons derived from stem cells display similar morphology nine months after transplantation into mouse cortex (Espuny-Camacho et al., 2013). However, despite the presence of synapses and action potentials, there is as yet no robust evidence for functioning neural networks with anatomically correct circuitry. This could be

because organoids lack the dorsoventral, anteroposterior and other axes found in embryonic brains, so although normal neuronal types differentiate efficiently in organoids, they are unlikely to be arranged in the appropriate anatomical locations relative to their prospective synaptic partners. Further, axonogenesis and synaptogenesis don't become prominent in human embryos until around 5-7 months (de Graaf-Peters and Hadders-Algra, 2006; Zhong et al., 2018), suggesting organoids are too immature. Some studies have characterized the electrophysiological properties of neurons in the human fetal cortex. One study using whole-cell patch clamping on slice culture of prefrontal cortex found no action potentials detected in neurons at GW23 and only detected APs at GW26 (Zhong et al., 2018), coinciding with the expression of axonogenesis genes during GW19-26.

While most organoid studies use transcriptomics to characterize their organoids, the presence of a transcript does not always indicate presence of functional protein. Expression of NMDA receptor subunit mRNA was detected in young human fetal neurons but its protein was not detected until >23 weeks (Eugenin et al., 2011). Therefore, expression of proteins in organoid neurons is likely a better marker of neuronal maturity.

Some electrophysiological properties, such as increased capacitance and sodium and potassium currents, mature over time in organoids, indicating neuronal maturation with age (Qian et al., 2016). Analysis of 10 month old organoids using multi-electrode arrays (MEA) showed consistently increased firing rates, burst frequency and synchrony indicating maturing neural networks. Notably, comparison of timing of electrical activity of pre-term human EEGs with the MEA recordings from cortical organoids showed organoids over 6 months old had high developmental age correlation, suggesting that the organoids follow an intrinsic developmental timeline (Trujillo et al., 2019). In a separate study, higher firing frequency was observed in thalamic organoids when they were fused to cortical organoids, suggesting that interactions between the organoids led to neuronal maturation (Xiang et al., 2019). Generating regionalized organoids may be limiting their maturation as extrinsic signals such as guidance molecules from neighbouring regions may be required for full maturation.

Human ESC-derived neurons transplanted into mouse brain integrated and matured over a period of several months, eventually generating action potentials similar to those seen in adult humans. Intrinsic properties improved over time; membrane potential hyperpolarization, increase in maximum sodium currents, increased firing rates, dendritic length and dendritic spine density.

Two-photon and calcium imaging experiments found an increase in calcium activity in human neurons during mouse visual stimulation indicating synaptic integration within the host cortex (Linaro et al., 2019).

These experiments show that stem cell derived neurons can mature to produce functionally similar neurons to those seen *in vivo* (Linaro et al., 2019; Trujillo et al., 2019) and can functionally integrate with local circuitry (Linaro et al., 2019). However, the process of maturation takes several months (6-10 months) (Linaro et al., 2019; Nicholas et al., 2013; Trujillo et al., 2019) following their own intrinsic developmental timeline independent of *in vitro* differentiation protocol or culture conditions (Linaro et al., 2019). Extrinsic signals can improve maturation (Xiang et al., 2019) indicating some culture conditions may be limiting cortical organoid maturation.

Despite their lack of maturation, the electrophysiological properties of cerebral organoids have been shown to model some aspects of disease phenotypes. For example, organoids harbouring mutations implicated in Alzheimer's disease (AD) displayed increased levels of neurotransmitter transporter proteins and increased AP firing rates. This hyperexcitability has also been seen in AD mouse models and human brains (Ghatak et al., 2019). Hyperexcitability was also seen in organoids modelling Angelman Syndrome (AS), a neurodevelopmental disorder partly characterized by seizures. When compared to wild type, AS-cortical organoids showed increased firing with neurons in some organoids showing synchronous firing (Sun et al., 2019). Changes in electrophysiological properties were also found in Timothy Syndrome cortical organoids which displayed abnormal calcium signalling (Birey et al., 2017) and idiopathic ASD-cerebral organoids which required a more hyperpolarized membrane potential to inactivate sodium channels (Mariani et al., 2015).

Organoids could well provide a useful tool for studying synaptic function in ASD but this will require improvements in differentiation protocols to allow normal maturation of neurons. This suggests that cerebral organoids, in their current form, are most appropriate for the study of earlier neurodevelopmental processes such as neurogenesis and cortical lamination rather than later processes such as synaptogenesis and neural circuit formation.

4. Challenges in using cerebral organoids to model ASDs

4.1 Differences between organoids and *in vivo* development

While there are many similarities between organoids and their *in vivo* counterparts, understanding the key differences can lead to improvements in the model. One major source of differences arises from the use of SMAD inhibition to direct stem cells toward neural fate. The human brain consists of more than just neural cells; non-ectodermal cells such as microglia, endothelial, blood cells and immune cells make up around 23-27% of fetal cortex at ages 15-37pcw (Fan et al., 2018; Nowakowski et al., 2017). Microglia make up ~20% of fetal cortex cells from scRNA-seq studies (Fan et al., 2018; Nowakowski et al., 2017), and may be involved in ASD pathogenesis by dysfunctional synaptic pruning (Koyama and Ikegaya, 2015). Neuroimmunology is a large and expanding field as such many labs are working on ways to incorporate an immune system within the organoid model (Ormel et al., 2018) reported that omitting SMAD inhibition gave rise to organoids containing all three germ layers, ultimately allowing microglia to develop within the organoid. Microglia can also become incorporated within organoids by co-culturing them with iPSC-derived microglia (Lin et al., 2018). A major difference from the *in vivo* situation is the lack of vascularization in organoids, commonly leading to a necrotic core at their centre. Most cells within day 44 cortical organoids were associated with glycolysis compared to just 2% in fetal cortex, suggesting that organoid cells are metabolically stressed (Amiri et al., 2008). This abnormal rate of glycolysis could mask effects caused by ASD-causing mutations as studies have shown elevated glycolysis as a candidate cause of ASD (Vallee and Vallee, 2018). Stress pathways were found to be upregulated in cortical organoids, irrespective of protocol, cell line or organoid age (Bhaduri et al., 2020). Furthermore, these high stress levels limit cell type specification of organoid cells (Bhadouri et al., 2020). Transcriptomic comparison between fetal tissues and organoids suggest that major differences are due to tissue culture environment and not due to differences in differentiation (Camp et al., 2015). To improve nutrient and oxygen diffusion, spinning bioreactors (Lancaster et al., 2013), shaking culture systems (Lancaster and Knoblich, 2014) and sliced organoids (Qian et al., 2020) have been used. Using an air-liquid interface culture improves organoid survival, morphology and axon outgrowth (Giandomenico et al., 2019). Lack of vascularisation could also be the cause of the most striking difference between organoids and their *in vivo* counterpart, their size, with organoids being considerably smaller, possibly suggesting that processes involved in cortical expansion (Box 2) are not well recapitulated in organoids. Recently, Qian et al., (2020) showed that sustained growth of organoids can be

maintained by slicing organoids open to improve diffusion. Progenitor zones of sliced organoids continued to expand over time and were packed with NPCs, IPCs and oRGs, which are key cell types involved in cortical expansion. Despite their higher proliferative capacity, organoids did not undergo gyrification, although this could be induced by embedding organoids in ECM (Qian et al., 2020).

4.2 Sources of Variation

One of the main challenges in using organoids is their variability. Different laboratories tend to have their own in-house protocol in the generation of organoids and protocols vary in their differentiation efficiency depending on which cell line is used (Bershteyn et al., 2017; Birey et al., 2017; Lancaster et al., 2013; Xiang et al., 2019). There are multiple sources of technical variation between organoid differentiation protocols, for example recombinant proteins used to promote differentiation, Matrigel, and serum all have some batch effects during their production. Ideally, it would be helpful to have a systematic analysis of all published protocols and an agreed standardized protocol to be used in organoid generation. Nonetheless, significant efforts have been made to tackle this issue. A recent study comparing individual organoids generated from different lines and different batches using single-cell RNA-seq analysis showed very low organoid-to-organoid variability when careful quality control checks such as checks for neural differentiation were used during the process of organoid differentiation (Velasco et al., 2019). A common issue when using PSCs in cell culture, particularly for prolonged culture or during stressful events such as gene editing, is their propensity to acquire genetic changes which may alter their growth, transcriptome and/or differentiation, ultimately confounding experimental results (Mills et al., 2013). PSCs appear to incur non-random genetic abnormalities, the most abundant being chromosomal duplications or deletions. These can be assayed by karyotyping or qPCR (Baker et al., 2016). More subtle mutations such as SNP or CNVs can also occur in PSCs with apparently normal karyotype thus SNP arrays and/or genome sequencing should be performed on all lines used during experiments (Mills et al., 2013). However, these techniques may miss some genetic alterations thus it is important to use multiple clonal lines for experiments.

5. Future perspectives

Organoids provide a powerful, amenable *in vitro* system to study normal embryonic human brain development and how it is disrupted in neurodevelopmental disorders. Advances in imaging technologies and gene editing have opened multiple avenues to dissect the roles of specific genes during human brain development using organoids. Using advanced imaging of cortical organoids, such as light-sheet live-cell imaging microscopy (Held et al., 2018), will allow scientists to track in real-time the high dynamic processes of human neurogenesis. However, there are still many challenges facing the use of cortical organoids for studying ASD. One of the most obvious is the lack of any behavioural readout from organoids. Currently, ASD is diagnosed solely on behavioural characteristics but organoid studies might elucidate cellular phenotypes/readouts that were missed from studies using other model systems, given that human neurodevelopment is still relatively understudied and there are important interspecies differences (Rakic, 2009). Recent advances in sequencing technology and data analysis have increased our ability to understand complex neurodevelopmental disorders such as ASD by finding converging pathways (Pinto et al., 2014; Wen et al., 2016). Coupling that with organoid technology, one can imagine sequencing and bioinformatic analyses of ASD patient-derived cerebral organoids might provide new biomarkers for diagnosing ASD, instead of relying on behavioural phenotypes. Synaptogenesis and neural circuit formation are significant points of convergence in ASD-risk genes. Electrophysiological studies of cerebral organoids are limited in their current state due to the long maturation time needed for circuit formation. This is not helped by culture conditions which appear to induce stress on the organoids, limiting or even regressing neuronal maturity (Bhaduri et al., 2020). Despite these limitations some cellular phenotypes of neurodevelopmental disorders have been modelled using cerebral organoids (Table 1). Moreover, now that we have identified stress factors that limit neuronal progenitor maturation in organoids (Bhaduri et al., 2020), we can proceed to improve organoid culture conditions, for example by redesigning cell culture media to better support neuronal maturation or the use of microfluidics to enhance nutrient intake. It is challenging to link genotype with behavioral phenotype in ASD as ASD linked mutations often show incomplete penetrance. For example, only 26% of individuals carrying the 16p11.2 duplication are diagnosed with ASD, and around 37% of carriers of the duplication have no psychiatric diagnosis (Niarchou et al., 2019). Furthermore, monozygotic twins only show a 70% ASD concordance rate (Folstein and Rosen-Sheidley, 2001). Stochastic events and/or

environmental differences during early brain development may make one twin more susceptible to the effects of the mutation, leading to a more severe phenotype. While we might consider the current variability in organoids to be a confounding element in interpreting experimental results, this variability could hold the key in understanding the stochasticity and/or environmental differences that we find in monozygotic twins. More systematic studies will have to be undertaken to explore this, but the advent of organoids allows us for the first time to test this hypothesis.

Although cerebral organoids contain most of the cell types found in the brain, some important cell types are absent, such as microglia, which normally originate outside the brain (Bhaduri et al., 2020; Tanaka et al., 2020). Therefore, it is still difficult to model cellular aspects of ASD pathologies such as neuroinflammation (which requires the presence of microglia) or white matter abnormalities. However, there are already efforts in increasing the complexity of the cerebral organoid culture system through the use of co-culture with other cell types such as microglia (can be used to model neuroinflammation) (Song et al., 2019) and co-culturing with different brain regions to study axon tracts (to study white matter abnormalities) (Birey et al., 2017; Cullen et al., 2019; Xiang et al., 2019).

Despite their current limitations, cerebral organoids provide us with a valuable additional tool to investigate the etiology of autism. As better high-throughput techniques such as scRNA-seq and osmFISH (Codeluppi et al., 2018) are developed, more information can be obtained from the limited human fetal and embryonic samples. This increased human data, coupled with cellular data from 2D and maturing human cerebral organoid and behavioural data from animal models, will together give us better tools to understand ASD.

Box 1

Embryonic development of the cerebral cortex: a primer

The development of the cerebral cortex can be divided into several stages (see Figure). **1.**

Patterning: The basic plan of the mammalian brain is laid out at the neural plate stage. At this stage, signaling centres that surround the neural plate produce signaling molecules including FGFs, BMPs and SHH which form a set of intersecting gradients across the neuroectoderm. Combinations of these signals are believed to confer specific regional fate on neural plate cells (Hebert and Fishell, 2008). Next, in the process of **neurulation (2)**, the edges of the neural plate fold towards each other and fuse together, thus forming the neural tube. At the earliest stages, the neural tube contains three primary brain vesicles, the prosencephalon (forebrain), mesencephalon (midbrain) and the rhombencephalon (hindbrain) (Ishikawa et al., 2012). The prosencephalon expands disproportionately, becoming larger than the other vesicles. Each vesicle subsequently gives rise to specific parts of the brain. For example, the prosencephalon will give rise to the cerebral cortex, ventral telencephalon, thalamus and hypothalamus (Wilson and Houart, 2004). In the early neural tube, neuroepithelial progenitor cells divide symmetrically at the ventricular edge, giving rise to two daughter progenitors. These proliferative divisions rapidly expand the pool of neural progenitors (Florio and Huttner, 2014; Rakic, 1974). Neuroepithelial progenitor cells subsequently transform to form other progenitor types, primarily radial glia. Radial glia may divide either symmetrically or asymmetrically, giving rise to a radial glial and a neuronal daughter, thus initiating the process of **neurogenesis (3)** in which cortical neurons are born. As development proceeds, an increasing proportion of radial glia divide asymmetrically, generating large numbers of neurons. Another important population of neural progenitors, known as intermediate progenitors or apical progenitors, are found in the subventricular zone (SVZ) (Florio and Huttner, 2014; Noctor et al., 2001; Noctor et al., 2004; Tan and Shi, 2013). Intermediate progenitors then continue to divide, making more neurons. Newborn neurons **migrate (4)** radially (indicated by black arrows) towards the outer (pial) edge of the embryonic cortex guided by a scaffold provided by radial glial fibers that project from the ventricular edge to the pial surface. Early-born neurons populate the deepest layers of the cortex. Later born neurons migrate past them, progressively building up the characteristic six-layered of the cortex, in the process of **lamination** (Rakic, 1974, 1988, 2009; Seto and Eiraku, 2019; Tan and Shi, 2013). Neurons in each cortical layer have distinct molecular signatures, associated with their

specific functions. Excitatory (glutamatergic) cortical neurons are generated in dorsal telencephalon, but inhibitory (GABAergic) cortical neurons are born in the ventral telencephalon, from where they migrate tangentially into the forming cortex. Once cortical neurons have migrated to their final destinations, they next form connections with their appropriate synaptic partners in the process of **synaptogenesis (5)**, the first step in neural circuit assembly. During synaptogenesis, cell adhesion molecules such as neuroligins and neuexins are recruited to the site of the future synapse where they form a bridge between the axon and dendrite. This initiates protein specialization to organize the active zone of the presynaptic terminal and the post synaptic density (PSD) over a period of hours to days (Okabe et al., 2001). During this process, scaffolding proteins such as membrane-associated guanylate kinases (MAGUKs), PSD95, and SHANK1 are recruited to the site of axo-dendritic membrane contact (Butz et al., 1998; Irie et al., 1997). Next is the process of **synapse stabilization (6)**. In rodents, thousands of synapses and dendritic spines per neuron are added in the period of 1 – 2 weeks of development but the majority of the synapses are removed or withdrawn and neuronal activity plays a key role in this (Gipson and Olive, 2017; Ozcan and Ozcan, 2018). Many of the proteins located in the developing PSD play a role in synapse stabilization as many were shown to regulate synapse number and size. Neuronal connections could be between neurons from other brain structures that are further away (long-ranged connectivity) or with neurons from the same region of the brain (local connectivity). These connections are not final as many connections are made throughout embryonic and early development of the brain which will then be refined later on in development as connections that are used more are strengthened (activity-based neural connections) while connections that are less used are pruned as describe in the process of synapse stabilization to establish mature neural circuits (Schuldiner and Yaron, 2015; Tierney and Nelson, 2009). Figure is modified, with permission, from Price et al., (2017).

Box 2

Cortical expansion in humans

There are significant differences in the number, types and behaviour of cortical neural progenitors between mouse and human (see Figure). In mouse, early neuroepithelial cells (NEC, coloured red) divide symmetrically, to increase the size of the neuroepithelium before transforming to become radial glial progenitors (RGC, coloured blue). RGCs and apical

intermediate progenitor cells (aIPCs, coloured green) initially self-renew but gradually begin to undergo asymmetric divisions to produce either an intermediate progenitor cell (IPC, coloured orange) or a neuron (grey) (Farkas and Huttner, 2008). Most IPCs divide only once, producing two neurons. Once the balance of radial glial divisions shifts from self-renewal toward differentiative divisions, final neuronal output becomes restricted. Newborn neurons migrate radially through the intermediate zone (IZ) and subplate (SP) before settling in the cortical plate (CP). Equivalents of each of these progenitor types are found in primates, where they show increased self-renewal which, along with an increased starting population, leads to a larger VZ and SVZ (Smart et al., 2002; Zecevic et al., 2005). Primates, including human, have two proliferative subventricular layers, the inner and outer subventricular zones (iSVZ and oSVZ respectively). The oSVZ contains a variety of highly proliferative progenitor cells including outer radial glia (oRG, coloured yellow) which give rise to large numbers of cortical neurons (Fietz et al., 2010; Hansen et al., 2010). oRG are abundant in human embryonic cortex, but extremely rare in mouse. Further, neurogenesis takes place for longer in human cortex compared to mouse, allowing more rounds of cell division. Overall, the combination of higher starting cell population, additional progenitor types, higher proliferative capacity of progenitors and longer time-window for neurogenesis have contributed to the large expansion of human cortex compared to mouse. Figure is modified, with permission, from Mason and Price (2016).

Declarations

Ethics approval and consent to participate: Not applicable

Consent for publication: Not applicable

Availability of data and materials: Not applicable

Competing interests: None

Funding: Work in the authors' laboratories is funded by the Simons Initiative for the Developing Brain SFARI – 429085.

Authors' contributions: WKC and RG drafted the manuscript. DJP and JOM edited and revised the draft manuscript. All authors read and approved the final manuscript

Acknowledgements: We are grateful to Dr Sally Till for her helpful comments on the manuscript.

Authors' information (optional): Not applicable

References

- Amiri, A., Coppola, G., Scuderi, S., Wu, F., Roychowdhury, T., Liu, F., Pochareddy, S., Shin, Y., Safi, A., Song, L., *et al.* (2018). Transcriptome and epigenome landscape of human cortical development modeled in organoids. *Science* 362.
- Anderlid, B.M., Schoumans, J., Anneren, G., Tapia-Paez, I., Dumanski, J., Blennow, E., and Nordenskjold, M. (2002). FISH-mapping of a 100-kb terminal 22q13 deletion. *Human genetics* 110, 439-443.
- Bagni, C., and Greenough, W.T. (2005). From mRNP trafficking to spine dysmorphogenesis: the roots of fragile X syndrome. *Nat Rev Neurosci* 6, 376-387.
- Baker, D., Hirst, A.J., Gokhale, P.J., Juarez, M.A., Williams, S., Wheeler, M., Bean, K., Allison, T.F., Moore, H.D., Andrews, P.W., and Barbaric, I. (2016). Detecting Genetic Mosaicism in Cultures of Human Pluripotent Stem Cells. *Stem cell reports* 7, 998-1012.
- Bedogni, F., Hodge, R.D., Elsen, G.E., Nelson, B.R., Daza, R.A., Beyer, R.P., Bammler, T.K., Rubenstein, J.L., and Hevner, R.F. (2010). Tbr1 regulates regional and laminar identity of postmitotic neurons in developing neocortex. *Proc Natl Acad Sci U S A* 107, 13129-13134.
- Bershteyn, M., Nowakowski, T.J., Pollen, A.A., Di Lullo, E., Nene, A., Wynshaw-Boris, A., and Kriegstein, A.R. (2017). Human iPSC-Derived Cerebral Organoids Model Cellular Features of Lissencephaly and Reveal Prolonged Mitosis of Outer Radial Glia. *Cell stem cell* 20, 435-449 e434.
- Bhaduri, A., Andrews, M.G., Leon, W.M., Jung, D., Shin, D., Allen, D., Jung, D., Schmunk, G., Haeussler, M., Salma, J., *et al.* (2020). Cell stress in cortical organoids impairs molecular subtype specification. *Nature* 578, 142-148.
- Bhandari, R., Paliwal, J., and Kuhad, A. (2020). Neuropsychopathology of Autism Spectrum Disorder: Complex Interplay of Genetic, Epigenetic, and Environmental Factors. *Advances in neurobiology* 24.
- Birey, F., Andersen, J., Makinson, C.D., Islam, S., Wei, W., Huber, N., Fan, H.C., Metzler, K.R.C., Panagiotakos, G., Thom, N., *et al.* (2017). Assembly of functionally integrated human forebrain spheroids. *Nature* 545, 54-59.
- Bonaglia, M.C., Giorda, R., Borgatti, R., Felisari, G., Gagliardi, C., Selicorni, A., and Zuffardi, O. (2001). Disruption of the ProSAP2 gene in a t(12;22)(q24.1;q13.3) is associated with the 22q13.3 deletion syndrome. *American journal of human genetics* 69, 261-268.
- Bozdagi, O., Sakurai, T., Papapetrou, D., Wang, X., Dickstein, D.L., Takahashi, N., Kajiwar, Y., Yang, M., Katz, A.M., Scattoni, M.L., *et al.* (2010). Haploinsufficiency of the autism-associated Shank3 gene leads to deficits in synaptic function, social interaction, and social communication. *Molecular autism* 1, 15.
- Brown, V., Jin, P., Ceman, S., Darnell, J.C., O'Donnell, W.T., Tenenbaum, S.A., Jin, X., Feng, Y., Wilkinson, K.D., Keene, J.D., *et al.* (2001). Microarray identification of FMRP-associated brain mRNAs and altered mRNA translational profiles in fragile X syndrome. *Cell* 107, 477-487.
- Butler, M.G., Dasouki, M.J., Zhou, X.P., Talebizadeh, Z., Brown, M., Takahashi, T.N., Miles, J.H., Wang, C.H., Stratton, R., Pilarski, R., and Eng, C. (2005). Subset of individuals with autism spectrum disorders and extreme macrocephaly associated with germline PTEN tumour suppressor gene mutations. *Journal of medical genetics* 42, 318-321.
- Butz, S., Okamoto, M., and Sudhof, T.C. (1998). A tripartite protein complex with the potential to couple synaptic vesicle exocytosis to cell adhesion in brain. *Cell* 94, 773-782.
- Camp, J.G., Badsha, F., Florio, M., Kanton, S., Gerber, T., Wilsch-Brauninger, M., Lewitus, E., Sykes, A., Hevers, W., Lancaster, M., *et al.* (2015). Human cerebral organoids recapitulate gene expression programs of fetal neocortex development. *Proc Natl Acad Sci U S A* 112, 15672-15677.
- Casanova, M.F. (2014). Autism as a sequence: from heterochronic germinal cell divisions to abnormalities of cell migration and cortical dysplasias. *Medical hypotheses* 83, 32-38.

709 CDC (2012). Prevalence of autism spectrum disorders- Autism and Developmental Disabilities
710 Monitoring Network, 14 sites, United States, 2008. Morbidity and mortality weekly report Surveillance
711 summaries (Washington, DC : 2002) 61, 1-19.

712 Chahrour, M., Jung, S.Y., Shaw, C., Zhou, X., Wong, S.T., Qin, J., and Zoghbi, H.Y. (2008). MeCP2, a key
713 contributor to neurological disease, activates and represses transcription. *Science* 320, 1224-1229.

714 Chahrour, M., and Zoghbi, H.Y. (2007). The story of Rett syndrome: from clinic to neurobiology. *Neuron*
715 56, 422-437.

716 Chao, H.T., Chen, H., Samaco, R.C., Xue, M., Chahrour, M., Yoo, J., Neul, J.L., Gong, S., Lu, H.C., Heintz, N.,
717 *et al.* (2010). Dysfunction in GABA signalling mediates autism-like stereotypies and Rett syndrome
718 phenotypes. *Nature* 468, 263-269.

719 Chen, R.Z., Akbarian, S., Tudor, M., and Jaenisch, R. (2001). Deficiency of methyl-CpG binding protein-2
720 in CNS neurons results in a Rett-like phenotype in mice. *Nature genetics* 27, 327-331.

721 Chen, Y., Huang, W.C., Sejourne, J., Clipperton-Allen, A.E., and Page, D.T. (2015). Pten Mutations Alter
722 Brain Growth Trajectory and Allocation of Cell Types through Elevated beta-Catenin Signaling. *J Neurosci*
723 35, 10252-10267.

724 Chow, M.L., Pramparo, T., Winn, M.E., Barnes, C.C., Li, H.R., Weiss, L., Fan, J.B., Murray, S., April, C.,
725 Belinson, H., *et al.* (2012). Age-dependent brain gene expression and copy number anomalies in autism
726 suggest distinct pathological processes at young versus mature ages. *PLoS Genet* 8, e1002592.

727 Codeluppi, S., Borm, L.E., Zeisel, A., La Manno, G., van Lunteren, J.A., Svensson, C.I., and Linnarsson, S.
728 (2018). Spatial organization of the somatosensory cortex revealed by osmFISH. *Nature methods* 15, 932-
729 935.

730 Comery, T.A., Harris, J.B., Willems, P.J., Oostra, B.A., Irwin, S.A., Weiler, I.J., and Greenough, W.T. (1997).
731 Abnormal dendritic spines in fragile X knockout mice: maturation and pruning deficits. *Proc Natl Acad Sci*
732 U S A 94, 5401-5404.

733 Conti, S., Condo, M., Posar, A., Mari, F., Resta, N., Renieri, A., Neri, I., Patrizi, A., and Parmeggiani, A.
734 (2012). Phosphatase and tensin homolog (PTEN) gene mutations and autism: literature review and a
735 case report of a patient with Cowden syndrome, autistic disorder, and epilepsy. *Journal of child*
736 *neurology* 27, 392-397.

737 Courchesne, V.H., G., and NE, L. (2020). Prenatal Origins of ASD: The When, What, and How of ASD
738 Development. *Trends in neurosciences* 43.

739 Courchesne, E., Carper, R., and Akshoomoff, N. (2003). Evidence of brain overgrowth in the first year of
740 life in autism. *Jama* 290, 337-344.

741 Courchesne, E., Mouton, P.R., Calhoun, M.E., Semendeferi, K., Ahrens-Barbeau, C., Hallet, M.J., Barnes,
742 C.C., and Pierce, K. (2011). Neuron number and size in prefrontal cortex of children with autism. *Jama*
743 306, 2001-2010.

744 Cullen, D.K., Gordián-Vélez, W.J., Struzyna, L.A., Jgamadze, D., Lim, J., Wofford, K.L., Browne, K.D., and
745 Chen, H.I. (2019). Bundled Three-Dimensional Human Axon Tracts Derived From Brain Organoids.
746 *iScience* 21.

747 Darnell, J.C., Jensen, K.B., Jin, P., Brown, V., Warren, S.T., and Darnell, R.B. (2001). Fragile X mental
748 retardation protein targets G quartet mRNAs important for neuronal function. *Cell* 107, 489-499.

749 de Graaf-Peters, V.B., and Hadders-Algra, M. (2006). Ontogeny of the human central nervous system:
750 what is happening when? *Early human development* 82, 257-266.

751 de la Torre-Ubieta, L., Won, H., Stein, J.L., and Geschwind, D.H. (2016). Advancing the understanding of
752 autism disease mechanisms through genetics. *Nature medicine* 22, 345-361.

753 De Rubeis, S., and Buxbaum, J.D. (2015). Genetics and genomics of autism spectrum disorder: embracing
754 complexity. *Human molecular genetics* 24, R24-31.

755 De Rubeis, S., He, X., Goldberg, A.P., Poultney, C.S., Samocha, K., Cicek, A.E., Kou, Y., Liu, L., Fromer, M.,
 756 Walker, S., *et al.* (2014). Synaptic, transcriptional and chromatin genes disrupted in autism. *Nature* 515,
 757 209-215.
 758 Dhar, S.U., del Gaudio, D., German, J.R., Peters, S.U., Ou, Z., Bader, P.I., Berg, J.S., Blazo, M., Brown,
 759 C.W., Graham, B.H., *et al.* (2010). 22q13.3 deletion syndrome: clinical and molecular analysis using array
 760 CGH. *Am J Med Genet A* 152A, 573-581.
 761 Durand, C.M., Betancur, C., Boeckers, T.M., Bockmann, J., Chaste, P., Fauchereau, F., Nygren, G., Rastam,
 762 M., Gillberg, I.C., Anckarsater, H., *et al.* (2007). Mutations in the gene encoding the synaptic scaffolding
 763 protein SHANK3 are associated with autism spectrum disorders. *Nature genetics* 39, 25-27.
 764 Espuny-Camacho, I., Michelsen, K.A., Gall, D., Linaro, D., Hasche, A., Bonnefont, J., Bali, C., Orduz, D.,
 765 Bilheu, A., Herpoel, A., *et al.* (2013). Pyramidal neurons derived from human pluripotent stem cells
 766 integrate efficiently into mouse brain circuits in vivo. *Neuron* 77, 440-456.
 767 Eugenin, E.A., King, J.E., Hazleton, J.E., Major, E.O., Bennett, M.V., Zukin, R.S., and Berman, J.W. (2011).
 768 Differences in NMDA receptor expression during human development determine the response of
 769 neurons to HIV-tat-mediated neurotoxicity. *Neurotoxicity research* 19, 138-148.
 770 Fan, X., Dong, J., Zhong, S., Wei, Y., Wu, Q., Yan, L., Yong, J., Sun, L., Wang, X., Zhao, Y., *et al.* (2018).
 771 Spatial transcriptomic survey of human embryonic cerebral cortex by single-cell RNA-seq analysis. *Cell*
 772 *research* 28, 730-745.
 773 Farkas, L.M., and Huttner, W.B. (2008). The cell biology of neural stem and progenitor cells and its
 774 significance for their proliferation versus differentiation during mammalian brain development. *Current*
 775 *opinion in cell biology* 20, 707-715.
 776 Fazel Darbandi, S., Robinson Schwartz, S.E., Qi, Q., Catta-Preta, R., Pai, E.L., Mandell, J.D., Everitt, A.,
 777 Rubin, A., Krasnoff, R.A., Katzman, S., *et al.* (2018). Neonatal Tbr1 Dosage Controls Cortical Layer 6
 778 Connectivity. *Neuron* 100, 831-845 e837.
 779 Feng, Y., Absher, D., Eberhart, D.E., Brown, V., Malter, H.E., and Warren, S.T. (1997). FMRP associates
 780 with polyribosomes as an mRNP, and the I304N mutation of severe fragile X syndrome abolishes this
 781 association. *Molecular cell* 1, 109-118.
 782 Fernandez, B.A., and Scherer, S.W. (2017). Syndromic autism spectrum disorders: moving from a
 783 clinically defined to a molecularly defined approach. *Dialogues in clinical neuroscience* 19, 353-371.
 784 Fietz, S.A., Kelava, I., Vogt, J., Wilsch-Brauninger, M., Stenzel, D., Fish, J.L., Corbeil, D., Riehn, A., Distler,
 785 W., Nitsch, R., and Huttner, W.B. (2010). OSVZ progenitors of human and ferret neocortex are epithelial-
 786 like and expand by integrin signaling. *Nat Neurosci* 13, 690-699.
 787 Fischbach, G.D., and Lord, C. (2010). The Simons Simplex Collection: a resource for identification of
 788 autism genetic risk factors. *Neuron* 68, 192-195.
 789 Florio, M., and Huttner, W.B. (2014). Neural progenitors, neurogenesis and the evolution of the
 790 neocortex. *Development* 141, 2182-2194.
 791 Folstein, S.E., and Rosen-Sheidley, B. (2001). Genetics of autism: complex aetiology for a heterogeneous
 792 disorder. *Nat Rev Genet* 2, 943-955.
 793 Gallagher, D., Voronova, A., Zander, M.A., Cancino, G.I., Bramall, A., Krause, M.P., Abad, C., Tekin, M.,
 794 Neilsen, P.M., Callen, D.F., *et al.* (2015). Ankrd11 is a chromatin regulator involved in autism that is
 795 essential for neural development. *Dev Cell* 32, 31-42.
 796 Gauthier, J., Champagne, N., Lafreniere, R.G., Xiong, L., Spiegelman, D., Brustein, E., Lapointe, M., Peng,
 797 H., Cote, M., Noreau, A., *et al.* (2010). De novo mutations in the gene encoding the synaptic scaffolding
 798 protein SHANK3 in patients ascertained for schizophrenia. *Proc Natl Acad Sci U S A* 107, 7863-7868.
 799 Ghatak, S., Dolatabadi, N., Trudler, D., Zhang, X., Wu, Y., Mohata, M., R, A., Talantova, M., and Lipton,
 800 S.A. (2019). Mechanisms of Hyperexcitability in Alzheimer's Disease hiPSC-derived Neurons and Cerebral
 801 Organoids vs Isogenic Controls. *eLife* 8.

Giandomenico, S.L., Mierau, S.B., Gibbons, G.M., Wenger, L.M.D., Masullo, L., Sit, T., Sutcliffe, M., Boulanger, J., Tripodi, M., Derivery, E., *et al.* (2019). Cerebral organoids at the air-liquid interface generate diverse nerve tracts with functional output. *Nat Neurosci* 22, 669-679.

Gilman, S.R., Iossifov, I., Levy, D., Ronemus, M., Wigler, M., and Vitkup, D. (2011). Rare de novo variants associated with autism implicate a large functional network of genes involved in formation and function of synapses. *Neuron* 70, 898-907.

Gipson, C.D., and Olive, M.F. (2017). Structural and functional plasticity of dendritic spines - root or result of behavior? *Genes, brain, and behavior* 16, 101-117.

Grove, J., Ripke, S., Als, T.D., Mattheisen, M., Walters, R.K., Won, H., Pallesen, J., Agerbo, E., Andreassen, O.A., Anney, R., *et al.* (2019). Identification of common genetic risk variants for autism spectrum disorder. *Nature Genetics* 51, 431-444.

Guang, S., Pang, N., Deng, X., Yang, L., He, F., Wu, L., Chen, C., Yin, F., and Peng, J. (2018). Synaptopathology Involved in Autism Spectrum Disorder. *Front Cell Neurosci* 12, 470.

Hagerman, R.J., Van Housen, K., Smith, A.C., and McGavran, L. (1984). Consideration of connective tissue dysfunction in the fragile X syndrome. *American journal of medical genetics* 17, 111-121.

Hansen, D.V., Lui, J.H., Parker, P.R., and Kriegstein, A.R. (2010). Neurogenic radial glia in the outer subventricular zone of human neocortex. *Nature* 464, 554-561.

Hazlett, H.C., Gu, H., Munsell, B.C., Kim, S.H., Styner, M., Wolff, J.J., Elison, J.T., Swanson, M.R., Zhu, H., Botteron, K.N., *et al.* (2017). Early brain development in infants at high risk for autism spectrum disorder. *Nature* 542, 348-351.

He, X., Sanders, S.J., Liu, L., De Rubeis, S., Lim, E.T., Sutcliffe, J.S., Schellenberg, G.D., Gibbs, R.A., Daly, M.J., Buxbaum, J.D., *et al.* (2013). Integrated model of de novo and inherited genetic variants yields greater power to identify risk genes. *PLoS Genet* 9, e1003671.

Hebert, J.M., and Fishell, G. (2008). The genetics of early telencephalon patterning: some assembly required. *Nat Rev Neurosci* 9, 678-685.

Held, M., Santeramo, I., Wilm, B., Murray, P., and Levy, R. (2018). Ex vivo live cell tracking in kidney organoids using light sheet fluorescence microscopy. *PLoS One* 13, e0199918.

Hendrich, B., and Bird, A. (1998). Identification and characterization of a family of mammalian methyl-CpG binding proteins. *Molecular and cellular biology* 18, 6538-6547.

Hevner, R.F., Shi, L., Justice, N., Hsueh, Y., Sheng, M., Smiga, S., Bulfone, A., Goffinet, A.M., Campagnoni, A.T., and Rubenstein, J.L. (2001). *Tbr1* regulates differentiation of the preplate and layer 6. *Neuron* 29, 353-366.

Hormozdiari, F., Penn, O., Borenstein, E., and Eichler, E.E. (2015). The discovery of integrated gene networks for autism and related disorders. *Genome research* 25, 142-154.

Huang, G., Chen, S., Chen, X., Zheng, J., Xu, Z., Doostparast Torshizi, A., Gong, S., Chen, Q., Ma, X., Yu, J., *et al.* (2019). Uncovering the Functional Link Between SHANK3 Deletions and Deficiency in Neurodevelopment Using iPSC-Derived Human Neurons. *Frontiers in neuroanatomy* 13, 23.

Hung, A.Y., Futai, K., Sala, C., Valtschanoff, J.G., Ryu, J., Woodworth, M.A., Kidd, F.L., Sung, C.C., Miyakawa, T., Bear, M.F., *et al.* (2008). Smaller dendritic spines, weaker synaptic transmission, but enhanced spatial learning in mice lacking Shank1. *J Neurosci* 28, 1697-1708.

Iossifov, I., O'Roak, B.J., Sanders, S.J., Ronemus, M., Krumm, N., Levy, D., Stessman, H.A., Witherspoon, K.T., Vives, L., Patterson, K.E., *et al.* (2014). The contribution of de novo coding mutations to autism spectrum disorder. *Nature* 515, 216-221.

Irie, M., Hata, Y., Takeuchi, M., Ichtchenko, K., Toyoda, A., Hirao, K., Takai, Y., Rosahl, T.W., and Sudhof, T.C. (1997). Binding of neuroligins to PSD-95. *Science* 277, 1511-1515.

Irwin, S.A., Galvez, R., and Greenough, W.T. (2000). Dendritic spine structural anomalies in fragile-X mental retardation syndrome. *Cerebral cortex* 10, 1038-1044.

Ishikawa, Y., Yamamoto, N., Yoshimoto, M., and Ito, H. (2012). The primary brain vesicles revisited: are the three primary vesicles (forebrain/midbrain/hindbrain) universal in vertebrates? *Brain Behav Evol* 79, 75-83.

Jordan, C., Li, H.H., Kwan, H.C., and Francke, U. (2007). Cerebellar gene expression profiles of mouse models for Rett syndrome reveal novel MeCP2 targets. *BMC medical genetics* 8, 36.

Kamiya, A., Kubo, K., Tomoda, T., Takaki, M., Youn, R., Ozeki, Y., Sawamura, N., Park, U., Kudo, C., Okawa, M., *et al.* (2005). A Schizophrenia-Associated Mutation of DISC1 Perturbs Cerebral Cortex Development. *Nature cell biology* 7.

Koyama, R., and Ikegaya, Y. (2015). Microglia in the pathogenesis of autism spectrum disorders. *Neurosci Res* 100, 1-5.

La Fata, G., Gartner, A., Dominguez-Iturza, N., Dresselaers, T., Dawitz, J., Poorthuis, R.B., Avena, M., Himmelreich, U., Meredith, R.M., Achsel, T., *et al.* (2014). FMRP regulates multipolar to bipolar transition affecting neuronal migration and cortical circuitry. *Nat Neurosci* 17, 1693-1700.

Laggerbauer, B., Ostareck, D., Keidel, E.M., Ostareck-Lederer, A., and Fischer, U. (2001). Evidence that fragile X mental retardation protein is a negative regulator of translation. *Human molecular genetics* 10, 329-338.

Lancaster, M.A., and Knoblich, J.A. (2014). Generation of cerebral organoids from human pluripotent stem cells. *Nature protocols* 9, 2329-2340.

Lancaster, M.A., Renner, M., Martin, C.A., Wenzel, D., Bicknell, L.S., Hurles, M.E., Homfray, T., Penninger, J.M., Jackson, A.P., and Knoblich, J.A. (2013). Cerebral organoids model human brain development and microcephaly. *Nature* 501, 373-379.

Lange, C., Huttner, W.B., and Calegari, F. (2009). Cdk4/cyclinD1 overexpression in neural stem cells shortens G1, delays neurogenesis, and promotes the generation and expansion of basal progenitors. *Cell stem cell* 5, 320-331.

Lewis, J.D., Meehan, R.R., Henzel, W.J., Maurer-Fogy, I., Jeppesen, P., Klein, F., and Bird, A. (1992). Purification, sequence, and cellular localization of a novel chromosomal protein that binds to methylated DNA. *Cell* 69, 905-914.

Li, Y., Muffat, J., Omer, A., Bosch, I., Lancaster, M.A., Sur, M., Gehrke, L., Knoblich, J.A., and Jaenisch, R. (2017). Induction of Expansion and Folding in Human Cerebral Organoids. *Cell stem cell* 20, 385-396 e383.

Li, Z., Zhang, Y., Ku, L., Wilkinson, K.D., Warren, S.T., and Feng, Y. (2001). The fragile X mental retardation protein inhibits translation via interacting with mRNA. *Nucleic Acids Res* 29, 2276-2283.

Lim, S., Naisbitt, S., Yoon, J., Hwang, J.I., Suh, P.G., Sheng, M., and Kim, E. (1999). Characterization of the Shank family of synaptic proteins. Multiple genes, alternative splicing, and differential expression in brain and development. *J Biol Chem* 274, 29510-29518.

Lin, Y.T., Seo, J., Gao, F., Feldman, H.M., Wen, H.L., Penney, J., Cam, H.P., Gjoneska, E., Raja, W.K., Cheng, J., *et al.* (2018). APOE4 Causes Widespread Molecular and Cellular Alterations Associated with Alzheimer's Disease Phenotypes in Human iPSC-Derived Brain Cell Types. *Neuron* 98, 1294.

Linaro, D., Vermaercke, B., Iwata, R., Ramaswamy, A., Davis, B.A., Boubakar, L., Libé-Philippot, B., Bilheu, A., Bruyne, L.D., Gall, D., *et al.* (2019). Xenotransplanted human cortical neurons reveal species-specific development and functional integration into mouse visual circuits. *bioRxiv*, 626218.

Luo, C., Lancaster, M.A., Castanon, R., Nery, J.R., Knoblich, J.A., and Ecker, J.R. (2016). Cerebral Organoids Recapitulate Epigenomic Signatures of the Human Fetal Brain. *Cell reports* 17, 3369-3384.

Manuel, M.N., Martynoga, B., Molinek, M.D., Quinn, J.C., Kroemmer, C., Mason, J.O., and Price, D.J. (2011). The transcription factor Foxg1 regulates telencephalic progenitor proliferation cell autonomously, in part by controlling Pax6 expression levels. *Neural development* 6, 9.

895 Marchetto, M.C., Carromeu, C., Acab, A., Yu, D., Yeo, G.W., Mu, Y., Chen, G., Gage, F.H., and Muotri, A.R.
 896 (2010). A model for neural development and treatment of Rett syndrome using human induced
 897 pluripotent stem cells. *Cell* 143, 527-539.
 898 Mariani, J., Coppola, G., Zhang, P., Abyzov, A., Provini, L., Tomasini, L., Amenduni, M., Szekely, A.,
 899 Palejev, D., Wilson, M., *et al.* (2015). FOXG1-Dependent Dysregulation of GABA/Glutamate Neuron
 900 Differentiation in Autism Spectrum Disorders. *Cell* 162, 375-390.
 901 Marshall, C.R., Noor, A., Vincent, J.B., Lionel, A.C., Feuk, L., Skaug, J., Shago, M., Moessner, R., Pinto, D.,
 902 Ren, Y., *et al.* (2008). Structural variation of chromosomes in autism spectrum disorder. *American*
 903 *journal of human genetics* 82, 477-488.
 904 Mellen, M., Ayata, P., Dewell, S., Kriaucionis, S., and Heintz, N. (2012). MeCP2 binds to 5hmC enriched
 905 within active genes and accessible chromatin in the nervous system. *Cell* 151, 1417-1430.
 906 Mills, J.A., Wang, K., Paluru, P., Ying, L., Lu, L., Galvao, A.M., Xu, D., Yao, Y., Sullivan, S.K., Sullivan, L.M.,
 907 *et al.* (2013). Clonal genetic and hematopoietic heterogeneity among human-induced pluripotent stem
 908 cell lines. *Blood* 122, 2047-2051.
 909 Miyashiro, K.Y., Beckel-Mitchener, A., Purk, T.P., Becker, K.G., Barret, T., Liu, L., Carbonetto, S., Weiler,
 910 I.J., Greenough, W.T., and Eberwine, J. (2003). RNA cargoes associating with FMRP reveal deficits in
 911 cellular functioning in *Fmr1* null mice. *Neuron* 37, 417-431.
 912 Miyoshi, G., and Fishell, G. (2012). Dynamic FoxG1 expression coordinates the integration of multipolar
 913 pyramidal neuron precursors into the cortical plate. *Neuron* 74, 1045-1058.
 914 Moessner, R., Marshall, C.R., Sutcliffe, J.S., Skaug, J., Pinto, D., Vincent, J., Zwaigenbaum, L., Fernandez,
 915 B., Roberts, W., Szatmari, P., and Scherer, S.W. (2007). Contribution of SHANK3 mutations to autism
 916 spectrum disorder. *American journal of human genetics* 81, 1289-1297.
 917 Moretti, P., Levenson, J.M., Battaglia, F., Atkinson, R., Teague, R., Antalffy, B., Armstrong, D., Arancio, O.,
 918 Sweatt, J.D., and Zoghbi, H.Y. (2006). Learning and memory and synaptic plasticity are impaired in a
 919 mouse model of Rett syndrome. *J Neurosci* 26, 319-327.
 920 Mullins, C., Fishell, G., and Tsien, R.W. (2016). Unifying Views of Autism Spectrum Disorders: A
 921 Consideration of Autoregulatory Feedback Loops. *Neuron* 89, 1131-1156.
 922 Niarchou, M., Chawner, S., Doherty, J.L., Maillard, A.M., Jacquemont, S., Chung, W.K., Green-Snyder, L.,
 923 Bernier, R.A., Goin-Kochel, R.P., Hanson, E., *et al.* (2019). Psychiatric disorders in children with 16p11.2
 924 deletion and duplication. *Translational psychiatry* 9, 8.
 925 Nicholas, C.R., Chen, J., Tang, Y., Southwell, D.G., Chalmers, N., Vogt, D., Arnold, C.M., Chen, Y.J., Stanley,
 926 E.G., Elefanty, A.G., *et al.* (2013). Functional maturation of hPSC-derived forebrain interneurons requires
 927 an extended timeline and mimics human neural development. *Cell stem cell* 12, 573-586.
 928 Noctor, S.C., Flint, A.C., Weissman, T.A., Dammerman, R.S., and Kriegstein, A.R. (2001). Neurons derived
 929 from radial glial cells establish radial units in neocortex. *Nature* 409, 714-720.
 930 Noctor, S.C., Martinez-Cerdeno, V., Ivic, L., and Kriegstein, A.R. (2004). Cortical neurons arise in
 931 symmetric and asymmetric division zones and migrate through specific phases. *Nat Neurosci* 7, 136-144.
 932 Nowakowski, T.J., Bhaduri, A., Pollen, A.A., Alvarado, B., Mostajo-Radji, M.A., Di Lullo, E., Haeussler, M.,
 933 Sandoval-Espinosa, C., Liu, S.J., Velmeshev, D., *et al.* (2017). Spatiotemporal gene expression trajectories
 934 reveal developmental hierarchies of the human cortex. *Science* 358, 1318-1323.
 935 Nuber, U.A., Kriaucionis, S., Roloff, T.C., Guy, J., Selfridge, J., Steinhoff, C., Schulz, R., Lipkowitz, B.,
 936 Ropers, H.H., Holmes, M.C., and Bird, A. (2005). Up-regulation of glucocorticoid-regulated genes in a
 937 mouse model of Rett syndrome. *Human molecular genetics* 14, 2247-2256.
 938 O'Donnell, W.T., and Warren, S.T. (2002). A decade of molecular studies of fragile X syndrome. *Annual*
 939 *review of neuroscience* 25, 315-338.
 940 Ohta, H., Nordahl, C.W., Iosif, A.M., Lee, A., Rogers, S., and Amaral, D.G. (2016). Increased Surface Area,
 941 but not Cortical Thickness, in a Subset of Young Boys With Autism Spectrum Disorder. *Autism research* :
 942 official journal of the International Society for Autism Research 9, 232-248.

943 Okabe, S., Miwa, A., and Okado, H. (2001). Spine formation and correlated assembly of presynaptic and
 944 postsynaptic molecules. *J Neurosci* 21, 6105-6114.
 945 Ormel, P.R., Vieira de Sa, R., van Bodegraven, E.J., Karst, H., Harschnitz, O., Sneeboer, M.A.M., Johansen,
 946 L.E., van Dijk, R.E., Scheefhals, N., Berdenis van Berlekom, A., *et al.* (2018). Microglia innately develop
 947 within cerebral organoids. *Nature communications* 9, 4167.
 948 Osterweil, E.K., Krueger, D.D., Reinhold, K., and Bear, M.F. (2010). Hypersensitivity to mGluR5 and
 949 ERK1/2 leads to excessive protein synthesis in the hippocampus of a mouse model of fragile X syndrome.
 950 *J Neurosci* 30, 15616-15627.
 951 Ozcan, A.S., and Ozcan, M.S. (2018). Population Dynamics and Long-Term Trajectory of Dendritic Spines.
 952 *Frontiers in synaptic neuroscience* 10, 25.
 953 Packer, A. (2016). Neocortical neurogenesis and the etiology of autism spectrum disorder. *Neuroscience*
 954 *and biobehavioral reviews* 64, 185-195.
 955 Pasca, A.M., Sloan, S.A., Clarke, L.E., Tian, Y., Makinson, C.D., Huber, N., Kim, C.H., Park, J.Y., O'Rourke,
 956 N.A., Nguyen, K.D., *et al.* (2015). Functional cortical neurons and astrocytes from human pluripotent
 957 stem cells in 3D culture. *Nature methods* 12, 671-678.
 958 Patterson, K.C., Hawkins, V.E., Arps, K.M., Mulkey, D.K., and Olsen, M.L. (2016). MeCP2 deficiency results
 959 in robust Rett-like behavioural and motor deficits in male and female rats. *Human molecular genetics*
 960 25, 5514-5515.
 961 Peca, J., Feliciano, C., Ting, J.T., Wang, W., Wells, M.F., Venkatraman, T.N., Lascola, C.D., Fu, Z., and Feng,
 962 G. (2011). Shank3 mutant mice display autistic-like behaviours and striatal dysfunction. *Nature* 472, 437-
 963 442.
 964 Pelka, G.J., Watson, C.M., Radziewicz, T., Hayward, M., Lahooti, H., Christodoulou, J., and Tam, P.P.
 965 (2006). Mecp2 deficiency is associated with learning and cognitive deficits and altered gene activity in
 966 the hippocampal region of mice. *Brain : a journal of neurology* 129, 887-898.
 967 Pilaz, L.J., McMahon, J.J., Miller, E.E., Lennox, A.L., Suzuki, A., Salmon, E., and Silver, D.L. (2016).
 968 Prolonged Mitosis of Neural Progenitors Alters Cell Fate in the Developing Brain. *Neuron* 89, 83-99.
 969 Pinto, D., Delaby, E., Merico, D., Barbosa, M., Merikangas, A., Klei, L., Thiruvahindrapuram, B., Xu, X.,
 970 Ziman, R., Wang, Z., *et al.* (2014). Convergence of genes and cellular pathways dysregulated in autism
 971 spectrum disorders. *American journal of human genetics* 94, 677-694.
 972 Pramparo, T., Lombardo, M.V., Campbell, K., Barnes, C.C., Marinero, S., Solso, S., Young, J., Mayo, M.,
 973 Dale, A., Ahrens-Barbeau, C., *et al.* (2015). Cell cycle networks link gene expression dysregulation,
 974 mutation, and brain maldevelopment in autistic toddlers. *Molecular systems biology* 11, 841.
 975 Qian, X., Jacob, F., Song, M.M., Nguyen, H.N., Song, H., and Ming, G.L. (2018). Generation of human
 976 brain region-specific organoids using a miniaturized spinning bioreactor. *Nature protocols* 13, 565-580.
 977 Qian, X., Nguyen, H.N., Song, M.M., Hadiono, C., Ogden, S.C., Hammack, C., Yao, B., Hamersky, G.R.,
 978 Jacob, F., Zhong, C., *et al.* (2016). Brain-Region-Specific Organoids Using Mini-bioreactors for Modeling
 979 ZIKV Exposure. *Cell* 165, 1238-1254.
 980 Qian, X., Su, Y., Adam, C., Deutschmann, A., Pather, S., Goldberg, E., Su, K., Li, S., Lu, L., Jacob, F., *et al.*
 981 (2020). Sliced Human Cortical Organoids for Modeling Distinct Cortical Layer Formation. *Cell stem cell*
 982 26.
 983 Qin, M., Kang, J., Burlin, T.V., Jiang, C., and Smith, C.B. (2005). Postadolescent changes in regional
 984 cerebral protein synthesis: an in vivo study in the FMR1 null mouse. *J Neurosci* 25, 5087-5095.
 985 Rakic, P. (1974). Neurons in rhesus monkey visual cortex: systematic relation between time of origin and
 986 eventual disposition. *Science* 183, 425-427.
 987 Rakic, P. (1988). Specification of cerebral cortical areas. *Science* 241, 170-176.
 988 Rakic, P. (2009). Evolution of the neocortex: a perspective from developmental biology. *Nat Rev*
 989 *Neurosci* 10, 724-735.

990 Ronemus, M., Iossifov, I., Levy, D., and Wigler, M. (2014). The role of de novo mutations in the genetics
 991 of autism spectrum disorders. *Nat Rev Genet* 15, 133-141.
 992 Roussignol, G., Ango, F., Romorini, S., Tu, J.C., Sala, C., Worley, P.F., Bockaert, J., and Fagni, L. (2005).
 993 Shank expression is sufficient to induce functional dendritic spine synapses in aspiny neurons. *J Neurosci*
 994 25, 3560-3570.
 995 Sacco, R., Gabriele, S., and Persico, A.M. (2015). Head circumference and brain size in autism spectrum
 996 disorder: A systematic review and meta-analysis. *Psychiatry research* 234, 239-251.
 997 Saito, T., Hanai, S., Takashima, S., Nakagawa, E., Okazaki, S., Inoue, T., Miyata, R., Hoshino, K., Akashi, T.,
 998 Sasaki, M., *et al.* (2011). Neocortical layer formation of human developing brains and lissencephalies:
 999 consideration of layer-specific marker expression. *Cerebral cortex* 21, 588-596.
 1000 Sala, C., Piech, V., Wilson, N.R., Passafaro, M., Liu, G., and Sheng, M. (2001). Regulation of dendritic
 1001 spine morphology and synaptic function by Shank and Homer. *Neuron* 31, 115-130.
 1002 Samaco, R.C., Mandel-Brehm, C., Chao, H.T., Ward, C.S., Fyffe-Maricich, S.L., Ren, J., Hyland, K., Thaller,
 1003 C., Maricich, S.M., Humphreys, P., *et al.* (2009). Loss of MeCP2 in aminergic neurons causes cell-
 1004 autonomous defects in neurotransmitter synthesis and specific behavioral abnormalities. *Proc Natl Acad*
 1005 *Sci U S A* 106, 21966-21971.
 1006 Schuldiner, O., and Yaron, A. (2015). Mechanisms of developmental neurite pruning. *Cellular and*
 1007 *molecular life sciences : CMLS* 72, 101-119.
 1008 Seto, Y., and Eiraku, M. (2019). Human brain development and its in vitro recapitulation. *Neurosci Res*
 1009 138, 33-42.
 1010 Shahbazian, M.D., Antalffy, B., Armstrong, D.L., and Zoghbi, H.Y. (2002). Insight into Rett syndrome:
 1011 MeCP2 levels display tissue- and cell-specific differences and correlate with neuronal maturation.
 1012 *Human molecular genetics* 11, 115-124.
 1013 Shen, W., Ba, R., Su, Y., Ni, Y., Chen, D., Xie, W., Pleasure, S.J., and Zhao, C. (2019). Foxg1 Regulates the
 1014 Postnatal Development of Cortical Interneurons. *Cerebral cortex* 29, 1547-1560.
 1015 Sirmaci, A., Spiliopoulos, M., Brancati, F., Powell, E., Duman, D., Abrams, A., Bademci, G., Agolini, E.,
 1016 Guo, S., Konuk, B., *et al.* (2011). Mutations in ANKRD11 cause KBG syndrome, characterized by
 1017 intellectual disability, skeletal malformations, and macrodontia. *American journal of human genetics* 89,
 1018 289-294.
 1019 Smart, I.H., Dehay, C., Giroud, P., Berland, M., and Kennedy, H. (2002). Unique morphological features of
 1020 the proliferative zones and postmitotic compartments of the neural epithelium giving rise to striate and
 1021 extrastriate cortex in the monkey. *Cerebral cortex* 12, 37-53.
 1022 Song, L., Yuan, X., Jones, Z., Vied, C., Miao, Y., Marzano, M., Hua, T., Sang, Q.X.A., Guan, J., Ma, T., *et al.*
 1023 (2019). Functionalization of Brain Region-specific Spheroids with Isogenic Microglia-like Cells. *Sci Rep* 9.
 1024 Sun, A., Yuan, Q., Fukuda, M., Yu, W., Yan, H., Lim, G., Nai, M., D'Agostino, G., Tran, H., Itahana, Y., *et al.*
 1025 (2019). Potassium Channel Dysfunction in Human Neuronal Models of Angelman Syndrome. *Science*
 1026 (New York, NY) 366.
 1027 Tan, X., and Shi, S.H. (2013). Neocortical neurogenesis and neuronal migration. *Wiley interdisciplinary*
 1028 *reviews Developmental biology* 2, 443-459.
 1029 Tanaka, Y., Cakir, B., Xiang, Y., Sullivan, G.J., and Park, I.H. (2020). Synthetic Analyses of Single-Cell
 1030 Transcriptomes From Multiple Brain Organoids and Fetal Brain. *Cell reports* 30.
 1031 Tierney, A.L., and Nelson, C.A., 3rd (2009). Brain Development and the Role of Experience in the Early
 1032 Years. *Zero to three* 30, 9-13.
 1033 Trujillo, C.A., Gao, R., Negraes, P.D., Gu, J., Buchanan, J., Preissl, S., Wang, A., Wu, W., Haddad, G.G.,
 1034 Chaim, I.A., *et al.* (2019). Complex Oscillatory Waves Emerging from Cortical Organoids Model Early
 1035 Human Brain Network Development. *Cell stem cell* 25, 558-569 e557.

Tudor, M., Akbarian, S., Chen, R.Z., and Jaenisch, R. (2002). Transcriptional profiling of a mouse model for Rett syndrome reveals subtle transcriptional changes in the brain. *Proc Natl Acad Sci U S A* 99, 15536-15541.

Vaccarino, F.M., Grigorenko, E.L., Smith, K.M., and Stevens, H.E. (2009). Regulation of cerebral cortical size and neuron number by fibroblast growth factors: implications for autism. *J Autism Dev Disord* 39, 511-520.

Vallee, A., and Vallee, J.N. (2018). Warburg effect hypothesis in autism Spectrum disorders. *Molecular brain* 11, 1.

Velasco, S., Kedaigle, A.J., Simmons, S.K., Nash, A., Rocha, M., Quadrato, G., Paulsen, B., Nguyen, L., Adiconis, X., Regev, A., *et al.* (2019). Individual brain organoids reproducibly form cell diversity of the human cerebral cortex. *Nature* 570, 523-527.

Wang, P., Lin, M., Pedrosa, E., Hrabovsky, A., Zhang, Z., Guo, W., Lachman, H.M., and Zheng, D. (2015). CRISPR/Cas9-mediated heterozygous knockout of the autism gene CHD8 and characterization of its transcriptional networks in neurodevelopment. *Molecular autism* 6, 55.

Wang, P., Mokhtari, R., Pedrosa, E., Kirschenbaum, M., Bayrak, C., Zheng, D., and Lachman, H.M. (2017). CRISPR/Cas9-mediated heterozygous knockout of the autism gene CHD8 and characterization of its transcriptional networks in cerebral organoids derived from iPS cells. *Molecular autism* 8, 11.

Weiler, I.J., Irwin, S.A., Klintsova, A.Y., Spencer, C.M., Brazelton, A.D., Miyashiro, K., Comery, T.A., Patel, B., Eberwine, J., and Greenough, W.T. (1997). Fragile X mental retardation protein is translated near synapses in response to neurotransmitter activation. *Proc Natl Acad Sci U S A* 94, 5395-5400.

Wen, Y., Alshikho, M.J., and Herbert, M.R. (2016). Pathway Network Analyses for Autism Reveal Multisystem Involvement, Major Overlaps with Other Diseases and Convergence upon MAPK and Calcium Signaling. *PLoS One* 11, e0153329.

Wilson, S.W., and Houart, C. (2004). Early steps in the development of the forebrain. *Dev Cell* 6, 167-181.

Wu, Y., Zhong, W., Cui, N., Johnson, C.M., Xing, H., Zhang, S., and Jiang, C. (2016). Characterization of Rett Syndrome-like phenotypes in Mecp2-knockout rats. *Journal of neurodevelopmental disorders* 8, 23.

Xiang, Y., Tanaka, Y., Cakir, B., Patterson, B., Kim, K.Y., Sun, P., Kang, Y.J., Zhong, M., Liu, X., Patra, P., *et al.* (2019). hESC-Derived Thalamic Organoids Form Reciprocal Projections When Fused with Cortical Organoids. *Cell stem cell* 24, 487-497 e487.

Yoon, S.J., Elahi, L.S., Pasca, A.M., Marton, R.M., Gordon, A., Revah, O., Miura, Y., Walczak, E.M., Holdgate, G.M., Fan, H.C., *et al.* (2019). Reliability of human cortical organoid generation. *Nature methods* 16, 75-78.

Zecevic, N., Chen, Y., and Filipovic, R. (2005). Contributions of cortical subventricular zone to the development of the human cerebral cortex. *J Comp Neurol* 491, 109-122.

Zhang, A., Yeung, P.L., Li, C.W., Tsai, S.C., Dinh, G.K., Wu, X., Li, H., and Chen, J.D. (2004). Identification of a novel family of ankyrin repeats containing cofactors for p160 nuclear receptor coactivators. *J Biol Chem* 279, 33799-33805.

Zhao, X., Leotta, A., Kustanovich, V., Lajonchere, C., Geschwind, D.H., Law, K., Law, P., Qiu, S., Lord, C., Sebat, J., *et al.* (2007). A unified genetic theory for sporadic and inherited autism. *Proc Natl Acad Sci U S A* 104, 12831-12836.

Zhong, S., Zhang, S., Fan, X., Wu, Q., Yan, L., Dong, J., Zhang, H., Li, L., Sun, L., Pan, N., *et al.* (2018). A single-cell RNA-seq survey of the developmental landscape of the human prefrontal cortex. *Nature* 555, 524-528.

Zoghbi, H.Y., and Bear, M.F. (2012). Synaptic dysfunction in neurodevelopmental disorders associated with autism and intellectual disabilities. *Cold Spring Harb Perspect Biol* 4.

1083

1084 **Figure Legend**

1085 **Figure 1** Sections of (A) 30 day, (B) 45 day and (C) 65 day old cerebral organoid grown from
1086 wild-type human iPSCs. (A) Immunostaining for progenitor marker PAX6 (red) and deep layer
1087 neuronal marker CTIP2 (green). (B) Immunostaining for intermediate progenitor marker TBR2
1088 (green), deep later neuronal marker TBR1 (red) and deep layer neuronal marker CTIP2 (white).
1089 (C) Immunostaining for progenitor marker PAX6 (yellow), deep layer neuronal marker CTIP2
1090 (red) and upper layer neuronal marker SATB2 (green). Progenitors are located interiorly, near
1091 the central lumen whereas differentiated neurons are located towards the outer edge of the
1092 organoid and multiple time-points show that neuronal layers are progressively established,
1093 resembling *in vivo* embryonic cortex. L = lumen, O = outer edge. Scale bar: 100µm.

Figure for Box 1





Figure for Box 2



1095